

## Properties of exponents

Let  $a$  and  $b$  be positive numbers with  $a \neq 1$ ,  $b \neq 1$  and let  $x$  and  $y$  be real numbers. Then:

A) Exponent Laws:

1.  $a^x a^y = a^{x+y}$

2.  $(a^x)^y = a^{xy}$

3.  $(ab)^x = a^x b^x$

4.  $\left(\frac{a}{b}\right)^x = \frac{a^x}{b^x}$

5.  $\frac{a^x}{a^y} = a^{x-y}$

## Properties of Logarithms

Let  $b$  be a positive real number with  $b \neq 1$ , and let  $x$  be any real number. Then:

1.  $\log_b(1) = 0$      i.e.  $b^0 = 1$

2.  $\log_b(b) = 1$      i.e.  $b^1 = b$

3.  $\log_b(b^x) = x$      i.e.  $b^x = b^x$

4.  $b^{\log_b(x)} = x$  if  $x > 0$

5.  $\log_b(MN) = \log_b(M) + \log_b(N)$

6.  $\log_b\left(\frac{M}{N}\right) = \log_b(M) - \log_b(N)$

7.  $\log_b(M^p) = p \log_b(M)$

8.  $\log_b(M) = \log_b(N) \iff M = N$

## The natural logarithm

This is the same as before but now we use base  $e$ . Since the log base  $e$  shows up so often we call it the **natural log**.

$$\log_e(x) = \ln(x)$$

We also use log base 10 very often so we abbreviate that as

$$\log_{10}(x) = \log(x).$$

Your calculator follows the same convention.

## Change of Base Formula

Let  $a, b, x$  be positive real numbers with  $a \neq 1, b \neq 1$ . Then

$$\log_a(x) = \frac{\log_b(x)}{\log_b(a)} \quad (\text{For any } b)$$

For the calculator you can use either base 10 or base  $e$ .

$$\log_a(x) = \frac{\log(x)}{\log(a)} \quad \text{OR} \quad \log_a(x) = \frac{\ln(x)}{\ln(a)}.$$